



er Alnus catkins edible; hard nut with green husk, pendulous; anti-inflammatory salicin in bark; impressed le Malus Rosaceae. ss=self-sterile; Sb=spur bearer; multi-purpose; K>=Keeps to; Root stocks M25	Italian Alder A. Cordata vigorous and erect. Unaffected by wind.
Cider [apple] 1-5=blossom, harvest months. Fruit colour and tree size.  Ashton Brown Jersey 4, Nov bittersweet orange striped fruit.	Hen's Turds Sep/Oct
Black Dabinett 4, late Nov, bittersweet, dark purple flush  Brown (or Bushy) French	Josey Sep/Oct, Bittersweet  Kingston Black 3, November, sweet  Michelin 3, late October, medium sweet.
Brown Snout 5, Sept-Nov, sweet yellow/green, small/med tree  Cambridge Queening N/a  Chaxhill Red Mid-May, mid-Oct,	Morgan Sweet 3, Aug ferments quickly. Sweet also early eater  Nine of Diamonds? has 10 red spots in flesh not 9 as may be expected
Chisel Jersey 4,Nov,bittersweet.Red flushed fruit on med tree. RIP Crimson King 3, Nov, bittersweet. Green/yellow scarlet. RIP	Overleaf N/a Pear Box ?, cider and dessert use, small, good cropper.
Dabinett 3, November, bittersweet  Dymock Red April/May, Sept/Oct, bittersweet RIP?  Fillbarrell 3, November, sweet	Plum Vitae ?, July / August. Both cider and dessert use.  Red Styre 3 N/a  Somerset Redstreak 3, November, sweet
Foxwhelp 3, November, sweet  Golden Spire May, Sept-Dec.Ex.Lancs 1850. multi. K>March	Stoke Red 3, November, sweet  Sweet Alford 3, November, sweet
Guillevic 3, November, sweet  Gypsy Red No details available  Harry Masters Jersey 3, Nov bittersweet. Yellow dark red med	Sweet Coppin 2, September-November, sweet  Tremlett's Bitter 3, November, sweet  Varlington Mill 3, November, sweet
Hastings 3, November, sweet  Crab [apple] native, produces clusters of white/pink flowers. Tolerates most soil types. Copes with	Yarlington Mill 3, November, sweet  a shade but gives lower yields
Dartmouth open branches, white flowers, Oct harvest. 12ft@10yrs Culinary [apple] - Dessert (D) & cooking (C) First number or month is time of blossom, second	John Downie lge gold fruit Sept good pollinator. 12ft@10yrs d is diameter of bush after 10 years, third is harvest time.
Adam's Pearmain 2, 10 Keeps for 2 months.  American Mother 5, 8 Late autumn. Slow irregular growth.  Ashmeads Kernel 2, ? Best flavoured. Pick Early Oct store until February.	Howgate Wonder May, early October. Firm, Cooks well.  Keswick Codling Polln grp B s-fertile, 8, Sept, tends biennial  Lemon Pippin mid-May, early Oct. UK poss. Norman origin.
Bloody Ploughman 4 s/s,May,Sep,k'p till Nov.Sb OK in cold  Bradleys Beauty Mid-May, early Oct. English known in C17	Newton Wonder mid May, early October. Firm, Cooks well.  Pitmaston Pineapple 5, Oct/Nov, keeps to Jan. Small, russet
Cats Head Mid-May, early Oct. English known in C17 RIP Christmas Pippin 3 s/s, May,mid Oct keeps till Feb.Sb.Found 2000	Queen Cox 3,12 Appleby Or,Kingston Bagpuize1950s Ribston Pippin 2 s/s, Oct/Nov keep till Feb.Sb. C, D & cider
Dumelow's Seedling aka Wellington May, early Oct K>May.  Ellison's Orange 4, 12 Sept. strongly scented when ripe.  George Cave 2, 12 Late summer. Often ready by end of July.	Scotch Bridget Oct/Nov.Triploid. stores 3 months. C, D if stored Warners King 2, 15 Oct-Jan. Triploid C Cooking to a froth.
Common <i>Fraxinus excelsior</i> to 30mnative, leaves fall early, seed left on tree good bird food. Stra Mountain <i>Sorbus aucuparia Joseph Rock</i> flowers white and hang in clusters around May-June for	
ch Fagus sylvatica monoecious, female borne in pairs, male catkins in spring shortly after leaves. Ferry Vaccinium myrtillus undershrub with sweet edible purple berries. Leaves help lower blood su	ıgar levels.
h, Chinese Redbark Betula albosinensis to 25M, W China. Peeling brown bark. Northern variety kthorn <i>Rhamnus cathartica</i> dioecious. Named for woody spine on end of each twig. Black fruit make bignonioides leaves secrete nectar. Not self-fertilizing. Fruit long beans w winged seeds. Chero	nildly poisonous. Tree most invasive!
ar of Lebanon, <i>Cedrus libani (atlantica) var 'silberspitz'</i> produces creamy white new growth tip stnut, Sweet or Spanish <i>Castanea sativa</i> To 25m. Pale yellow catkins July, fruits November. Often	s. These stand out against the blue-green, inner needles on older branches en coppiced, almost rot proof, makes excellent fencing. First arrived at the time of the Armada. Needs a well-drain
ress, Lawson's 'Grayswood Feather' is a compact conifer of columnar habit, with soft, semi-juver pinnate leaves w.leaflets w. serrated margins white or cream-colored hermaphrodite flowers. All grays are the colored hermaphrodite flowers.	reen parts of plant poisonous.
Common <i>S.nigra</i> Berry extract effective against flu.  Brown Turkey' <i>Ficus carica</i> 'Fruit' is enclosed, lined on the inside with tiny flowers. Pollinated by the sof <i>Pinaceae</i> . Leaves attached singly with cup-like bases. Cones upright like related Cedrus, disingly with cup-like bases.	
Nordman A. nordmanniana 60m Ice Age refugia on S&E Black Sea coast.	Korean A. koreana 10-18m Blue cones.  Edible and appetite suppressants also used as a sweetener, sometimes used to flavour fish soup. Evergreen thick to
shaped leaves shiny dark green upper surface. <b>A. Sweet Liquidambar styraciflua</b> < 35m 2m dia. Leaves like maple tho gloss leathery. Resin used to	
<b>Thorn </b> <i>Crataegus</i> 5–15m, small pome "haw" fruit with 1-5 pyrenes like "stones" of plums. Thorns <b>Common </b> <i>C.monogyna</i> aka May,Quick- or White-thorn, Motherdie <b>el, Common </b> <i>Coryllus avellana</i> flowers in spring b4 leaves monoecious catkins, males yellow, fema	Glastonbury C.oxyacantha var.praecox Joseph of Aramathea's staff?
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and t	nd fall in late autumn. cones are erect, small, green or purple, ripening brown 5-8 months after pollination; norther grain for modelling, carving it small, round, ribbed cream nutlet somewhat toxic to bumblebees.
per <i>Juniperis communis</i> foliage of the juvenile needle-like type, unlike pine. Dioecious, male and the <i>Larix decidua</i> 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small cones with short bracts.  Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little a Broad-leaved Lime <i>T.platyphyllos</i> yellow-white flowers (make lime tea) pollinated by bees, fruit B Silver lime <i>T.tomentosa</i> flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is C Small-leaved lime <i>T.cordata</i> flowers green-yellow, hermaphrodite, edible in salads when young the <i>Acer</i> Of the family <i>Sapindaceae</i> . Sturdy widely branched deciduous tree with handsome palmate leaver the family <i>Sapindaceae</i> . Sturdy widely branched deciduous tree with handsome palmate leaver the family <i>Sapindaceae</i> . Sturdy widely branched deciduous tree with handsome palmate leaver the family sapindaceae. Sturdy widely branched deciduous tree with handsome palmate leaver the family sapindaceae. Sturdy widely branched deciduous tree with handsome palmate leaver the family sapindaceae. Sturdy widely branched deciduous tree with handsome palmate leaver the family sapindaceae. Sturdy widely branched deciduous tree with handsome palmate leaver the family sapindaceae. Sturdy widely branched deciduous tree with handsome palmate leaver the family sapindaceae. Sturdy widely branched deciduous tree with handsome palmate leaver the family sapindaceae.	r-like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour. female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  Independent on the second of the seco
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and the Larix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small cones with short bracts.  Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little a Broad-leaved Lime T.platyphyllos yellow-white flowers (make lime tea) pollinated by bees, fruit B Silver lime T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is C Small-leaved lime T.cordata flowers green-yellow, hermaphrodite, edible in salads when young the Acer Of the family Sapindaceae. Sturdy widely branched deciduous tree with handsome palmate to Amur A.ginnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange of Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from 'Ariadne' after grower's grand-daughter. Deeply cut leaves vividly marked with pink and white. 'Butterfly' light green leaves flecked with cream. In spring often edged with pink, in autumn the dissectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit	r-like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour.  female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  Ind fall in late autumn. cones are erect, small, green or purple, ripening brown 5-8 months after pollination; norther  grain for modelling, carving  it small, round, ribbed cream nutlet  is somewhat toxic to bumblebees.  it, much visited by bees. Fruit oval  eaves, mostly tolerating limestone soils and wind.  to red in autumn.  In drying winds and sun. To 5m.  Winter bark attractive. 2m arching habit  e cream areas turn deep scarlet
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and for the Larix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small cones with short bracts.  Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little a Broad-leaved Lime T.platyphyllos yellow-white flowers (make lime tea) pollinated by bees, fruit a Billian T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is a Small-leaved lime T.cordata flowers green-yellow, hermaphrodite, edible in salads when young the Acer Of the family Sapindaceae. Sturdy widely branched deciduous tree with handsome palmate to Amur A.ginnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange of Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from 'Ariadne' after grower's grand-daughter. Deeply cut leaves vividly marked with pink and white. 'Butterfly' light green leaves flecked with cream. In spring often edged with pink, in autumn the	r-like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour.  female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  Ind fall in late autumn. cones are erect, small, green or purple, ripening brown 5-8 months after pollination; norther  grain for modelling, carving  it small, round, ribbed cream nutlet  s somewhat toxic to bumblebees.  g, much visited by bees. Fruit oval  eaves, mostly tolerating limestone soils and wind.  to red in autumn.  In drying winds and sun. To 5m.  Winter bark attractive. 2m arching habit  te cream areas turn deep scarlet  urgined red, strong colour from good sun  mental effect / to control size
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and the Larix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small cones with short bracts.  Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little a Broad-leaved Lime T.platyphyllos yellow-white flowers (make lime tea) pollinated by bees, fruing Silver lime T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is a Silver lime T.cordata flowers green-yellow, hermaphrodite, edible in salads when young the Acer Of the family Sapindaceae. Sturdy widely branched deciduous tree with handsome palmate leads and the Amur A.ginnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange of Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from 'Ariadne' after grower's grand-daughter. Deeply cut leaves vividly marked with pink and white. 'Butterfly' light green leaves flecked with cream. In spring often edged with pink, in autumn the dissectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit  Maple A.shirasawanum 'Aureum' slow growing small tree. Yellow leaves in spring, in autumn matelly is Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for ornativity of Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for ornativity of Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for ornativity of Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for ornativity of Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for ornativity of Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for ornativity of Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for ornativity of Gold A.negundo golden leaves, long, golden flowe	-like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour. female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  Independent of all in late autumn. cones are erect, small, green or purple, ripening brown 5-8 months after pollination; norther grain for modelling, carving it small, round, ribbed cream nutlet is somewhat toxic to bumblebees.  It is much visited by bees. Fruit oval eaves, mostly tolerating limestone soils and wind. It is red in autumn. In drying winds and sun. To 5m.  Winter bark attractive. 2m arching habit eream areas turn deep scarlet  I rigined red, strong colour from good sun mental effect / to control size utumn. Red flowers in spring flowers on different racemes.  I spring.
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and the Larix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small cones with short bracts.  Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little a Broad-leaved Lime T.platyphyllos yellow-white flowers (make lime tea) pollinated by bees, frui B Silver lime T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is C Small-leaved lime T.cordata flowers green-yellow, hermaphrodite, edible in salads when young the Acer Of the family Sapindaceae. Sturdy widely branched deciduous tree with handsome palmate leading A.ginnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange of Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from 'Ariadne' after grower's grand-daughter. Deeply cut leaves vividly marked with pink and white. 'Butterfly' light green leaves flecked with cream. In spring often edged with pink, in autumn the dissectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit  Maple A.shirasawanum 'Aureum' slow growing small tree. Yellow leaves in spring, in autumn matelly's Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for ornar Nikko A.maximowiczianum grey-black peeling bark and trifoliate, green leaves that turn red in automakebark A.capillipes 10–15m often multiple trunks. Bark olive green, grey stripes. Male/female Sugar A.saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers early Sycamore A.pseudoplatanus 'brilliantissimum' slow growing intense salmon pink leaves, later yelloular Common Nottingham Mespilus germanica Russett apple-like fruit requires bletting to eat (slare Common Nottingham Mespilus germanica Russett apple-like fruit requires bletting to eat (slare Common Nottingham Mespilus germanica)	-like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour. female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  Independent of all in late autumn. cones are erect, small, green or purple, ripening brown 5-8 months after pollination; norther grain for modelling, carving it small, round, ribbed cream nutlet is somewhat toxic to bumblebees.  It is much visited by bees. Fruit oval eaves, mostly tolerating limestone soils and wind. It is red in autumn. In drying winds and sun. To 5m.  Winter bark attractive. 2m arching habit eream areas turn deep scarlet  I rigined red, strong colour from good sun mental effect / to control size utumn. Red flowers in spring flowers on different racemes.  I spring.
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and the Larix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small cones with short bracts.  Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little a Broad-leaved Lime T.platyphyllos yellow-white flowers (make lime tea) pollinated by bees, frui B Silver lime T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is C Small-leaved lime T.cordata flowers green-yellow, hermaphrodite, edible in salads when young the Acer Of the family Sapindaceae. Sturdy widely branched deciduous tree with handsome palmate leading A.ginnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange of Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from 'Ariadne' after grower's grand-daughter. Deeply cut leaves vividly marked with pink and white. 'Butterfly' light green leaves flecked with cream. In spring often edged with pink, in autumn the dissectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit  Maple A.shirasawanum 'Aureum' slow growing small tree. Yellow leaves in spring, in autumn matelly's Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for ornar Nikko A.maximowiczianum grey-black peeling bark and trifoliate, green leaves that turn red in automakebark A.capillipes 10–15m often multiple trunks. Bark olive green, grey stripes. Male/female Sugar A.saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers early Sycamore A.pseudoplatanus 'brilliantissimum' slow growing intense salmon pink leaves, later yelloular Common Nottingham Mespilus germanica Russett apple-like fruit requires bletting to eat (slare Common Nottingham Mespilus germanica Russett apple-like fruit requires bletting to eat (slare Common Nottingham Mespilus germanica)	-like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour. female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  Independent and fall in late autumn. cones are erect, small, green or purple, ripening brown 5-8 months after pollination; norther grain for modelling, carving  It small, round, ribbed cream nutlet  It small, round, ribbed cream nutlet  It somewhat toxic to bumblebees.  It much visited by bees. Fruit oval  It eaves, mostly tolerating limestone soils and wind.  It to red in autumn.  In drying winds and sun. To 5m.  Winter bark attractive. 2m arching habit  It e cream areas turn deep scarlet  It right red, strong colour from good sun  Immental effect / to control size  It utumn. Red flowers in spring  If owers on different racemes.  It yspring.  It green or purple, ripening brown 5-8 months after pollination; norther  It is mall, round, ribbed cream nutlet  It is mall, green or purple, ripening brown 5-8 months after pollination; norther  It is mall, green or purple, ripening brown 5-8 months after pollination; norther  It is mall, green or purple, ripening brown 5-8 months after pollination; norther  It is mall, green or purple, ripening brown 5-8 months after pollination; norther  It is mall, green or purple, ripening brown 5-8 months after pollination; norther  It is mall, green or purple, ripening brown 5-8 months after pollination; norther  It is mall, green or purple, ripening brown 5-8 months after pollination; norther  It is mall, green or purple, ripening brown 5-8 months after pollination; norther  It is mall, green or purple, ripening brown 5-8 months after pollination; norther  It is mall, green or purple, ripening brown 5-8 months after pollination; norther  It is mall, green or purple, ripening brown 5-8 months after pollination; norther  It is mall, green or purple, ripening brown 5-8 months after pollination; norther  It is mall, green or purple, ripening brown 5-8 months after pollination; norther
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and the Larix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small cones with short bracts.  Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little A Broad-leaved Lime T.platyphyllos yellow-white flowers (make lime tea) pollinated by bees, fruit B Silver lime T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is C Small-leaved lime T.cordata flowers green-yellow, hermaphrodite, edible in salads when young the Acer Of the family Sapindaceae. Sturdy widely branched deciduous tree with handsome palmate lead Amur A.ginnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from Ariadne' after grower's grand-daughter. Deeply cut leaves vividly marked with pink and white. Butterfly' light green leaves fleeked with cream. In spring often edged with pink, in autumn the dissectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit Maple A.shirasawanum 'Aureum' slow growing small tree. Yellow leaves in spring, in autumn maxedly's Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for orna Nikko A.maximowiczianum grey-black peeling bark and trifoliate, green leaves that turn red in an Snakebark A.capillipes 10–15m often multiple trunks. Bark olive green, grey stripes. Male/female Sugar A.saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers early Sycamore A.pseudoplatanus 'brilliantissimum' slow growing intense salmon pink leaves, later yellow and the purple fruit. SW Asia, 154 pairs of chromosom Quercus beech family, Fagaceae. Spirally arranged leaves, acorns contain tannic acid. Catkins and sn Pin Quercus petraea 20–40M white oak stalked leaves, stalkless (sessile) acorns. petraea means "Bin Quercus petraea	-like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour. female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  In different for modelling, carving it small, green or purple, ripening brown 5-8 months after pollination; norther grain for modelling, carving it small, round, ribbed cream nutlet is somewhat toxic to bumblebees.  In much visited by bees. Fruit oval eaves, mostly tolerating limestone soils and wind.  In ord in autumn.  In drying winds and sun. To 5m.  Winter bark attractive. 2m arching habit e cream areas turn deep scarlet  In urgined red, strong colour from good sun mental effect / to control size utumn. Red flowers in spring flowers on different racemes.  In y spring.  In wy spring.  In decay and ferment). Ornamental and hardy, slow and graceful in growth. White flowers in June.  In the season of the same for the same fruit & green parts intoxicating hall female flowers in spring.  In different flowers in spring.
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and the Larix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small cones with short bracts.  Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little A Broad-leaved Lime T.platyphyllos yellow-white flowers (make lime tea) pollinated by bees, frui B Silver lime T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is E Small-leaved lime T.cordata flowers green-yellow, hermaphrodite, edible in salads when young lea Acer Of the family Sapindaceae. Sturdy widely branched deciduous tree with handsome palmate lea Acer Of the family Sapindaceae. Sturdy widely branched deciduous tree with handsome palmate lea Apalmatum small slow growing, in autumn gold orange and red. Underplant to shelter from Analmatum small slow growing, in autumn gold orange and red. Underplant to shelter from Analmatum factor and the sectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit Maple A.shirasawanum 'Aureum' slow growing small tree. Yellow leaves in spring, in autumn ma Kelly's Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for orna Nikko A.maximowiczianum grey-black peeling bark and trifoliate, green leaves that turn red in an Snakebark A.capillipes 10–15m often multiple trunks. Bark olive green, grey stripes. Male/female Sugar A.saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers early Sycamore A.pseudoplatanus 'brilliantissimum' slow growing intense salmon pink leaves, later yellowers plack Morus nigra 10–15m. Self-fertile, dark purple fruit. SW Asia, 154 pairs of chromosom Quercus beech family, Fagaceae. Spirally arranged leaves, acorns contain tannic acid. Catkins and sne Pin Quercus petraea 20–40M white oak stalked leaves, stalkless (sessile) acorns. petraea means "ian Ironwood Parrotia persica to 30m smooth bark flakes and peels. Le	-like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour. female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  In dial in late autumn. cones are erect, small, green or purple, ripening brown 5-8 months after pollination; norther grain for modelling, carving  It small, round, ribbed cream nutlet  It somewhat toxic to bumblebees.  It small, round, ribbed cream nutlet  It swows mostly tolerating limestone soils and wind.  It ored in autumn.  In drying winds and sun. To 5m.  Winter bark attractive. 2m arching habit  It cream areas turn deep scarlet  Interpret of the control size  Intumn. Red flowers in spring  Inflowers on different racemes.  It spring.  In we green. Monoecious.  It starting to decay and ferment). Ornamental and hardy, slow and graceful in growth. White flowers in June.  It is spring from the spring and flowers ones droop  In for ones droop  In for ones droop  In for motelling the red stained, juicy and musky QA  Sanguinole 2, Aug/Sep. Small, russet, red blush & spotted. Flesh red stained, juicy and musky QA
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and the Laris decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are so have small cones with short bracts.  Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little A Broad-leaved Lime T.platyphyllos yellow-white flowers (make lime tea) pollinated by bees, frui B Silver lime T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is C Small-leaved lime T.cordata flowers green-yellow, hermaphrodite, edible in salads when young let Acer Of the family Sapindaevae. Sturdy widely branched deciduous tree with handsome palmate to Amur A.ginnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange is Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from "Ariadne" after grower's grand-daughter. Deeply cut leaves vividly marked with pink, in autumn the dissectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit Maple A.shirasawanum "Aureum" slow growing small tree. Yellow leaves in spring, in autumn ma Kelly's Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for orna Nitko A.maximowiczianum grey-black pecling bark and trifoliate, green leaves that turn red in at Snakebark A.capillipes 10–15m often multiple trunks. Bark olive green, grey stripes. Male/female Eugar A.saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers early Sycamore A.pseudoplatanus brilliantissimum' slow growing intense salmon pink leaves, later yelle lar Common Nottingham Mespilus germanica Russett apple-like fruit requires bletting to eat (septery Black Morus nigra 10–15m. Self-fertile, dark purple fruit. SW Asia, 154 pairs of chromosom Quercus petraea 20–40M white oak stalked leaves, stalkless (sessile) acorns. petraea men seian Ironwood Parrotia persica to 30m smooth bark flakes and peels. Leaves ovoid	Like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour. formale on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen. In dial in late autumn. cones are erect, small, green or purple, ripening brown 5-8 months after pollination; northe grain for modelling, carving its small, round, ribbed cream nutlet is somewhat toxic to bumblebees.  15. much visited by bees. Fruit oval eaves, mostly tolerating limestone soils and wind. It ored in autumn. In drying winds and sun. To 5m.  16. Winter bark attractive. 2m arching habit e cream areas turn deep scarlet eream areas turn deep scarlet utumn. Red flowers in spring.  17. flowers on different racemes.  18. y spring.  18. was green. Monoecious.  18. starting to decay and ferment). Ornamental and hardy, slow and graceful in growth. White flowers in June.  18. sensaches pruned after leaves fall can be used to make baskets. White sap in raw fruit & green parts intoxicating interest after for furniture making.  18. lilen A  18. Sanguinole 2, Aug/Sep. Small, russet, red blush & spotted. Flesh red stained, juicy and musky QA  18. ment Nellis  18. saller, Pyrus Communis, Wick Court  19. Judge Amphlett Rec, 1, late September, 1 week, med A, low T, avg Q
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and th Laris decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are se have small cones with short bracts.  Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little A Broad-leaved Lime T.platyphyllos yellow-white flowers (make lime tea) pollinated by bees, frui B Silver lime T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is C Small-leaved lime T.cordata flowers green-yellow, hermaphrodite, edible in salads when young le Acer Of the family Sapindaceae. Sturdy widely branched deciduous tree with handsome palmate be Amur A.ginnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange is Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from Irriadne' after grower's grand-daughter. Deeply cut leaves vividly marked with pink and white. Butterfly' light green leaves flecked with cream. In spring often edged with pink in autumn the dissectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit Maple A.shirasawanum 'Aureum' slow growing small tree. Yellow leaves in spring, in autumn ma Kelly's Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for orna Nikko A.maximowiczianum grey-black peeling bark and trifoliate, green leaves that turn red in at Snakebark A.capillipes 10-15m often multiple trunks. Bark olive green, grey stripes. Male/female Bugar A.saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers early Sycamore A.pseudoplatanus' brilliantissimum' slow growing intense salmon pink leaves, later yelle lar Common Nottingham Mespilus germanica Russett apple-like fruit requires bletting to eat (s beerry Black Morus nigra 10-15m. Self-fertile, dark purple fruit. SW Asia, 154 pairs of chromosom Quercus beech family, Fagaceae. Spirally arranged leaves, acoms contain tannic	Like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour.  female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  and fall in late autumn. cones are erect, small, green or purple, ripening brown 5-8 months after pollination; northe grain for modelling, carving it small, round, ribbed cream mudet somewhat toxic to bumblebees.  fa much visited by bees. Fruit oval eaves, mostly tolerating limestone soils and wind.  to red in autumn.  In drying winds and sun. To 5m.  Winter bark attractive. 2m arching habit e cream areas turn deep scarlet  regined red, strong colour from good sun mental effect / to control size  utumn. Red flowers in spring flowers on different racemes.  y spring.  yo green. Monoecious.  starting to decay and ferment). Ornamental and hardy, slow and graceful in growth. White flowers in June.  leas. Branches pruned after leaves fall can be used to make baskets. White sap in raw fruit & green parts intoxicating mall female flowers in spring.  izontal, lower ones droop  of rocky places?".  not purple in autumn  at after for furniture making.  lilien A  Sanguinole 2, Aug/Sep. Small, russet, red blush & spotted. Flesh red stained, juicy and musky QA  Thompson's Sep, store to Oct or Nov. Golden skin, parch of russet, Flesh juicy with rich perfume  Williams Bon Chretien 2,12 Aug/Sep.Eat fresh Ripe yellow QA  Winter Nellis  saller, Pyrus Communis, Wick Court  Judge Amphlett Rec, 1, late September, 1 week, med A, low T, avg Q  Mooreroft 2, late Sept, M 2d, med A, med T, astringent good Q  Newbridge 3, 1-3wk Oct, M<1vk, low A, Low T, avg Q
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and the Lark decidua 15-50 melaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small cones with short bracts.  E Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little A Broad-leaved Lime T.platyphyllos yellow-white flowers (make lime tea) pollinated by bees, frui B Silver lime T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectat is C Small-leaved lime T.cordata flowers green-yellow, hermaphrodite, edible in salads when young the Acer Of the family Sapiudaceae. Sturdy widely branched deciduous tree with handsome palmate leading the Acer of the family Sapiudaceae. Sturdy widely branched deciduous tree with handsome palmate leading the Acer of the family Sapiudaceae. Sturdy widely branched deciduous tree with handsome palmate leading the Acer of the family Sapiudaceae. Sturdy widely branched deciduous tree with handsome palmate leading to the Acer of the family Sapiudaceae. Sturdy widely branched deciduous tree with handsome palmate leading the Acer of the family Sapiudaceae. Sturdy widely branched deciduous tree with handsome palmate leading the Acer of the family Sapiudaceae. Sturdy widely branched deciduous tree with handsome palmate leading the Acer of the family Sapiudaceae. Sturdy widely branched deciduous tree with handsome palmate leading the Acer of the family Sapiudaceae. Sturdy widely branched deciduous tree with handsome palmate leading the Acer of the family Sapiudaceae. Sturdy widely branched deciduous tree with handsome palmate leaves branched family Sapiudaceae. Sturdy widely branched deciduous tree with handsome palmate leaves tree family sapiudaceae. Sturdy widely branched deciduous tree with handsome palmate leaves. In the Acer of the	Like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour.  fernale on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  do fall in late autumn. cones are erect, small, green or purple, ripening brown 5-8 months after pollination; norther  grain for modelling, carving  tt small, round, ribbed cream nutlet  s somewhat toxic to bumblebees.  s, much visited by bees. Fruit oval  eaves, mostly tolerating limestone soils and wind.  to red in autumn.  ndrying winds and sun. To 5m.  Winter bark attractive. 2m arching habit  e cream areas turn deep scarlet  urgined red, strong colour from good sun  mental effect / to control size  utumn. Red flowers in spring  flowers on different racemes.  y spring.  wy green. Monoecious.  bus Branches pruned after leaves fall can be used to make baskets. White sap in raw fruit & green parts intoxicating hall female flowers in spring,  iziontal, lower ones droop  of rocky places?  and purple in autumn  a tafer for furniture making,  alien A  Sanguinole 2, Aug/Sep. Small, russet, red blush & spotted. Flesh red stained, juicy and musky QA  rdens  Thompson's Sep, store to Oct or Nov. Golden skin, patch of russet, Flesh juicy with rich perfume  Williams Bon Chretien 2,12 Aug/Sep.Eat fresh Ripe yellow QA  Winter Nellis  saller, Pyrus Communis, Wick Court  Judge Amphlett Ree, 1, late September, 1 week, med A, low T, avg Q  Moorcroft 2, late Sept, M 2d, med A, med T, astringent good Q  Newbridge 3, 1-3wk Oct, M<1wk, low A, Low T, avg Q  Oldfield 2, Oct, 3-6 wks, med-hi A, med T, astringent good Q  Red Pear 5, Oct, ≥3 wks, lowA, lowT, avg Q good Q  Bed Pear 5, Oct, >3 wks, lowA, lowT, avg Q good Q  Red Pear 5, Oct, >3 wks, lowA, lowT, avg Q good Q  Snake Pole From Oxenton, very old general purpose
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and th Laix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are so have small cones with short bracts.  e Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little A Broad-leaved Lime T.platyphyllos yellow-white flowers (make lime tea) pollinated by bees, fruit B sliver lime T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is Silver lime T.tomentosa flowers green-yellow, hermaphrodite, edible in salads when young let Acer Of the family Sapindaeae. Sturdy widely branched deciduous tree with handsome palmate hanur A.ginnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from Ariadne' after grower's grand-daughter. Deeply cut leaves vividly marked with pink, in autumn the dissectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit Maple A.shirasawanum 'Aureum' slow growing small tree. Yellow leaves in spring, in autumn makelly's Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for ornat winklko A.maximowiczianum grey-black peeling bark and trifoliate, green leaves that turn red in at sinakebark A.capillipes 10–15m often multiple trunks. Bark olive green, grey stripes. Male female Bugar A.saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers carb systemor A.pseudoplatamus 'brilliantssimum' slow growing intense salmon pink leaves, later yelle lar Common Nottingham Mespilus germanica Russett apple-like fruit requires bletting to eat (s betty Black Morus nigra 10–15m. Self-fertile, dark purple fruit. SW Asia, 154 pairs of chromosom Quercus beech family, Fugaraea. Spirally arranged leaves, acorns contain tannic acid. Catkins and sn Prus deeply rooted, branching freely if cut, grows large to a great age, rivall	-like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour.  female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  In fall in late autumn. comes are creek, small, green or purple, ripening brown 5-8 months after pollination; norther  grain for modelling, carving  it small, round, ribbed cream nutlet  somewhat toxic to bumblebees.  , much visited by bees. Fruit oval  caves, mostly tolerating limestone soils and wind.  to red in autumn.  In drving winds and sun. To 5m.  Winter bark attractive. 2m arching habit  e cream areas turn deep scarlet  regined red, strong colour from good sun  mental effect / to control size  utumn. Red flowers in spring  flowers on different racemes.  y spring.  we green. Monoecious.  starting to decay and ferment). Ornamental and hardy, slow and graceful in growth. White flowers in June.  tess. Branches pruned after leaves fall can be used to make baskets. White sap in raw fruit & green parts intoxicating  nall female flowers in spring.  izontal, lower ones droop  for rocky places?  not purple in autumn  nt after for furniture making.  lilen A   Sanguinole 2, Aug/Sep. Small, russet, red blush & spotted. Flesh red stained, juicy and musky QA  rdens Thompson's Sep, store to Oct or Nov. Golden skin, patch of russet, Flesh juicy with rich perfume  Williams Bon Chretien 2,12 Aug/Sep.Eat fresh Ripe yellow QA  Winter Nellis   saller, Pyrus Communis, Wick Court  Judge Amphlett Rec, 1, late September, 1 week, med A, low T, avg Q  Oldfield 2, Oct, 3-6 wks, med-hi A, med T, avg to good Q  Red Pear 5, Oct, 2-3 wks, low A, low T, avg Q  Oldfield 2, Oct, 3-6 wks, med-hi A, med T, avg to good Q  Snake Pole From Oxenton, very old general purpose  Speart From Arlingham  Stoneyway N/a
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and the Lark decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are so have small cones with short bracts.  Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little A Broad-leaved Lime T.platpyllos yellow-white flowers (make lime tea) pollinated by bees, fruit a Bisilver lime T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is Silver lime T.tomentosa flowers pale yellow, hermaphrodite, edible in salads when young lee Acer Of the family Sapindaeaae. Sturdy widely branched deciduous tree with handsome palmate be Amur A.ginnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from Ariadne and the same state of the same shape and the dissectum Deceply cut leaves flecked with cream. In spring often edged with pink, in autumn the dissectum Deceply cut leaves flecked with cream. In spring often edged with pink, in autumn makelly's Gold A.negundo golden leaves, long, golden flowers in June. Prune hard in spring for ornat sinklko A.naximoviczianum grey-black peeling bark and trifoliate, green leaves that turn red in at Snakebark A.capillipes 10–15m often multiple trunks. Bark olive green, grey stripes. Male/female Surgar A.saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers early Sycamore A.pseudoplatanum Srvilliantsismum' slow growing intense salmon pink leaves, later yellow a proving intense salmon pink leaves, later yellow proving intense salmon pink leaves, later yellow a proving intense salmon pink leaves, later yellow and proving pink leaves the proving pink leaves the proving pink leaves the proving pink lea	Like fruits dark red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour.  female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  In fall in late autumn. cones are crect, small, green or purple, ripening brown 5-8 months after pollination; norther grain for modelling, carving  it small, round, ribbed cream nutlet  somewhat toxic to bumblebees.  a much visited by bees. Fruit oval  caves, mostly tolerating limestone soils and wind.  to red in autumn.  In drying winds and sun. To 5m.  Winter bark attractive. 2m arching habit  e ream areas turn deep scarlet  argined red, strong colour from good sun  mental effect / to control size  utumn. Red flowers in spring  flowers on different racenes.  y spring.  yw green. Monoecious.  starting to decay and ferment). Omamental and hardy, slow and graceful in growth. White flowers in June.  ses. Branches pruned after leaves fall can be used to make baskets. White sap in raw fruit & green parts intoxicating  hall female flowers in spring.  itsonal, lower ones droop  of rocky places.'  and purple in autumn  a rater for furniture making.  dien A   Sanguinole 2, Aug/Sep. Small, russet, red blush & spotted. Flesh red stained, juicy and musky QA  red  Winter Nellis   Sanguinole 3, 1-3wk Oct, M-Slwk, klow A, Low T, avg Q  Moorcroft 2, late Sept, M 2d, med A, med T, astringent good Q  Newbridge 3, 1-3wk Oct, M-Slwk, klow A, Low T, avg Q   Moldfield 2, Oct, 3-6 wks, med-hi A, med T, avg to good Q   Red Pear 5, Oct, >3 wks, lowA, lowT, avg Q   Snake Pole From Oxenton, very old general purpose  Spear Hrom Arlingham  Stoncyway N/a   Sweet Huffcap=Helens Early 3, Sept, 1 wk, med A, low—med T, avg Q   Thom 2, late Sept, 1 week, med A, low T, avg to good Q   From 2, late Sept, 1 week, med A, low T, avg to good Q   Red Pear 5, Oct, >3 wks, lowA, lowT, avg to good Q   Red Pear 5, Oct, >3 wks, lowA, lowT, avg to good Q   Red Pear 5, Oct, >6 wks, med-hi A, med T, avg to good Q   Red Pear 5, Oct, >6 wks, med-hi A, med T, avg
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and the Lark decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are se have small cones with short bracts.  Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little A Broad-leaved Lime T.platyphyllos yellow-white flowers (make lime tea) pollinated by bees, fruit as Silver lime T.tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is 2 Small-leaved lime T.cordata flowers green-yellow, hermaphrodite, edible in salads when young let Acer Of the family. Sapindaceae. Sturdy widely branched deciduous tree with handsome palmate lea Manua Aginnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange:  Amur A.ginnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange:  Ariadne' after grower's grand-daughter. Deeply cut leaves vividly marked with pink and white.  Butterfly light green leaves flecked with cream. In spring often edged with pink, in autumn the dissectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit  Maple A.shirasawanum Mureum' slow growing small tree. Yellow leaves in spring, in autumn makelly's Gold A.negundo golden leaves, long, golden flowers in lune. Prune hard in spring for orna Nikko A.maximowiczianum grey-black pecling bark and trifoliate, green leaves that turn red in anakebark A.capillipes 10-15m often multiple trunks. Bark olive green, grey stripes. Male/female Sugar A.saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers early sycamor A.pseudoplatanus brilliantsismum' slow growing intense salmon pink leaves, later yellow are manned to the proper strip of the server of the serve	Like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour.  female on separate plants, wind pollinated. Beryl-like seeds, taking 18 months to ripen.  In diall in late autumn. comes are erect, small, green or purple, ripening brown 5-8 months after pollination; norther  grain for modelling, carving  it small, round, ribbed cream nutlet  somewhat toxic to bumblebees.  , much visited by bees. Fruit oval  caves, mostly tolerating limestone soils and wind.  to red in autumn.  In drying winds and sun. 10-5m.  Winter bark attractive. 2m arching habit  c cream areas turn deep searlet   winter bark attractive. 2m good sun  mental effect / to control size   utumn. Red flowers in spring  flowers on different racemes.  y spring.   wing green. Monoecious.   starting to decay and ferment). Ornamental and hardy, slow and graceful in growth. White flowers in June.   sea. Branches pruned after leaves fall can be used to make baskets. White sap in raw fruit & green parts intoxicating  hall female flowers in spring.   izontal, lower ones droop   'of rocky places''.   Ind purple in autumn  at after for furniture making.  liken A   Sanguinole 2, Aug/Sep. Small, russet, red blush & spotted. Flesh red stained, juicy and musky QA   ridens Thompson's Sep, store to Oct or Nov. Golden skin, patch of russet, Flesh juicy with rich perfume   Williams Bon Chretine 2,12 Aug/Sep.Eat fresh Ripe yellow QA   Moreroft 2, late Sept, M 2d, med A, med T, astringent good Q   Newbridge 3, 1-3wk Oct, M-N will, Mo W, Low T, avg Q   Oldfield 2, Oct, ≥3 wks, low A, low'T, avg to good Q   Red Pear 3, Oct, ≥3 wks, low A, low'T, avg Q   Snake Pole From Oxenton, very old general purpose   Speart From Arlingham  Stoneyway N/a   Sweet Huffeap=Helens Barly 3, Sept, 1 wk, med A, low—med T, avg Q
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioceious, male and ih Larix decidua 15:50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are se have small cones with short brates.  Titia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little A Broad-leaved Lime T. Platyphylos yellow. white flowers (make lime tea) pollinated by bees, fini is sliver lime T. tomentosa flowers pale yellow, hermaphrodite, edible in salads when young de Acer Of the family Sapindaeau. Sturdy widely branched deciduous tree with handsome palmate by Amur A. ginnala bark thin, dull grav brown, smooth then fissured with age. Leaves brilliant orange i Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from Maple A.palmatum small slow growing rowing small tree. Yellow leaves in spring, in autumn the dissectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit Maple A.shirasawanum 'Aureum' slow growing small tree. Yellow leaves in spring, in autumn the Maple A.shirasawanum 'Aureum' slow growing small tree. Yellow leaves that turn red in as Snakebark A.capillipes 10–15m often multiple trunks. Bark olive green, grey stripes. Male/female Sugar A.saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers earb Sycamore A.pseudoplatanus 'brilliantissimum' slow growing intense salmon pink leaves, later yellow lar Common Nottingham Mespilus germanica Russett apple-like fruit requires bletting to cat (see her Maple A. Gramater and A. Sep-Oct A. St. Justice and A. Saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers carb Sycamore A. pseudoplatanus' brilliantissimum' slow growing intense salmon pink leaves, later yellow lar Common Nottingham Mespilus germanic	Like Fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour. female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  Ind fall in late autumn. cones are erect, small, green or purple, ripening brown 5-8 months after pollination; northe grain for modelling, carving  is small, round, ribbed cream authout the some street of the property of the property of the street of the pollination; northed as somewhat toxic to bumblebees.  In much visited by bees. Fruit oval  caves, mostly tolerating limestone soils and wind.  To red in autumn.  In dring winds and sun. To 5m.  Writer bark attractive. 2m arching habit comments that the comment of the property
per Junipetis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and in harix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small cones with short bracts.  2 Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little A Broad-leaved Lime T. Jaraphyrlios vellow-white flowers (make lime tea) pollinated by bees, fruig 3 Silver lime T. tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is 3 Silver lime T. tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is 3 Silver lime T. tomentosa flowers green-yellow, hermaphrodite, edible in salads when young let Acer Of the family Saphiakaea. Sturdy widely branched deciduous tree with handsome palmate be Amur A. gimnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange i Maple A.palmatum small slow growing, in autumn gold orange and red. Underplant to shelter fror I Ariadne' after grower's grand-daughter. Deeply cut leaves vividiy marked with pink, in autumn the dissectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit Maple A.shirasawanum Yaurcum 'slow growing small tree. Yellow leaves in spring, in autumn ma Kelly's Gold A.ncgundo golden leaves, long, golden flowers in lune. Prune hard in spring for orna silkko A.maximowicziaum grey-black pecking bark and tirioliate; green leaves that turn red in a sinakebark A.capillipes 10–15m often multiple trunks. Bark olive green, grey stripes. Male/female Sugar A.saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers early Sycamore A.pseudoplatams 'brilliantissium' slow growing intense salmon pink leaves, later yello lar Common Nottingham Mespilus germanica Russett apple-like fruit requires bletting to cat (see the growing silvers) and the surface of the silvers of chromosom Quercus beech family, Fugayaea. Sprilaly arranged leaves, acrons contain tamic aid clarentae red a. Prus	Like Fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour. female on separate plants, wind pollinated. Berry-like seeds, taking 18 months to ripen.  Ind fall in late autumn. comes are crect, small, green or purple, ripening brown 5-8 months after pollination; norther prain for modelling, carving it small, round, ribbed cream nudet is somewhat toxic to bumblebees.  In much vasited by bees, Fruit oval caves, mostly tolerating limestone soils and wind: to red in autumn.  In driving winds and sun. To 5m.  Winter bark attractive. 2m arching habit ceram areas turn deep seadet urgined red, strong colour from good sun mental effect. It to control size uturns. Red flowers in spring flowers on different racemes.  In spring.  In spring monoccious.  Itatring to decay and ferment). Ornamental and hardy, slow and praceful in growth. White flowers in lune.  Itatring to decay and ferment). Ornamental and hardy, slow and praceful in growth. White flowers in lune.  Itatring to decay and ferment). Ornamental and hardy, slow and praceful in growth. White flowers in lune.  Itatring to decay and ferment). Ornamental and hardy, slow and praceful in growth. White flowers in lune.  Itatring to decay and ferment). Ornamental and hardy, slow and praceful in growth. White flowers in lune.  Itatring to decay and ferment). Ornamental and hardy, slow and praceful in growth. White flowers in lune.  Itatring to decay and ferment). Ornamental and hardy, slow and praceful in growth. White flowers in lune.  Itatring to decay and ferment making.  Itatring to decay and ferment
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and In Larix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow at es have small cones with short bracts.  2 Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little A Broad-leaved Lime T. Palarphylios yellow: white flowers (make lime tea) pollinated by bees, fruil 3 Bilver lime T. tomentosa flowers: pale yellow, hermaphrodite, pollinated by honeybees. Nectar is C. Small-leaved lime T. Cordata flowers: pale yellow, hermaphrodite, pollinated by honeybees. Nectar is C. Small-leaved lime T. Cordata flowers: preen-yellow, hermaphrodite, colible in salads when young lie Acer Of the family Supradaeaa. Sturdy widely branched deciduous tree with handsome palmate to Manut A. Azimala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange in Maple A. Apalmatum small slow growing, in autumn gold orange and red. Underplant to shelter from Ariadne' after grower's grand-daughter. Deeply cut leaves vividly marked with pink and white. Butterfly light green leaves the handsome small tree. Yellow leaves in spring, in autumn the dissectum Deeply cut leaves. In high with broad, drooping habit.  Maple A. shirasawanum 'Aureum' slow growing small tree. Yellow leaves in spring, in autumn makelly's Gold A. negundo golden leaves, long, golden flowers in lune. Prune hard in spring for orna Nikko A. maximoviczianum grey-black pecling bark and trifoliate, green leaves that turn red in a linakebark A. explitipes 10–15m soften multiple trunks. Bark offwe green, grey stripes. Male/female Sugar A. saccharimum 45m. Leaves bright yellow to orange to fluorescent red-orange. Plowers early Sugar A. saccharimum 45m. Leaves bright yellow to orange to fluorescent red-orange. Plowers early Sugar A. saccharimum 45m. Leaves bright green leaves, starkets green leaves that turn red in a binakebark A. explitipes 10–15m soften multiple trunks. Bark offwe gree	Like fruits dark red when young, but become dark purple when ripe, edible and have an apple and sweet flavour. female on separate plants, wind polilinated. Bern-like seeds, taking 18 months to ripen.  Ind fall in late autumn, cones are erect, small, green or purple, ripening brown 5-8 months after pollination; northe grain for modelling, carving  It small, round, ribbed cream mutett  somewhat toxic to bumblebees.  In und, ribbed cream mutett  somewhat toxic to bumblebees.  In und visited by bees. Fruit oval  cases, mostly tolerating limestone soils and wind.  In red in autumn.  In drying winds and sun. To 5m.  Whiter bark attractive. 2m arching habit  e cream areas turn deep scarlet  rigined red, strong colour from good sun  mental effect / to control size  unumn. Red flowers in spring.  flowers on different racenes.  y spring.  we green. Monoecious.  starting to deepa and ferment). Ornamental and hardy, slow and graceful in growth. White flowers in lune.  seas Branches pruned after leaves fall can be used to make baskets. White sap in raw fruit & green parts intoxicating in purple in autumn  it after for furniture making.  Idien A  Sanguinote 2, Aug/Sep. Small, russet, red blush & spotted. Flesh red stained, juicy and musky QA  ridens  Sanguinote 2, Jug/Sep. Small, russet, red blush & spotted. Flesh red stained, juicy and musky QA  ridens  Sanguinote 2, Jug/Sep. Small, russet, red blush & spotted. Flesh red stained, juicy and musky QA  ridens  Milliams Bon Chretien 2, 12 Aug/Sep. Eat fresh Ripe vellow QA  Williams Bon Chretien 2, 12 Aug/Sep. Eat fresh Ripe vellow QA  Williams Bon Chretien 2, 12 Aug/Sep. Eat fresh Ripe vellow QA  Williams Bon Chretien 2, 12 Aug/Sep. Jay, Jay, Jay, Jay, Jay, Jay, Q  Oldfield 2, Oct., 3-6 wks, med-bi A, med T, awg to good Q  Newbridge 3, 1-3 wk Oct, M<1 wk, low A, Low T, avg Q  Oldfield 2, Oct., 3-6 wks, med-bi A, jow T, avg to good Q  New Hilliams Bon Chretien 2, 12 Aug/Sep. Jay, Jay, Jay, Jay, Jay, Jay, Jay, Jay,
per Juniperis communis foliage of the iuvenile needle-like type, unlike pine. Dioecious, male and in Larix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small cones with short bracts.  2 Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little ABroad-leaved Lime T. Pathyphyllos yellow-white flowers (make lime tea) pollinated by bees, fruit as its sturdy trunk, branches divide yellow, hermaphrodite, pollinated by honeybees. Nectar is 3 Silver lime T. tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is 3 Silver lime T. tomentosa flowers pale yellow, hermaphrodite, pollinated by honeybees. Nectar is 3 Silver lime T. tomentosa flowers pale yellow, hermaphrodite, celible in salads when young let Acer Of the family Japhinakara. Surdy widely branched deciduous tree with handsome palmate by Amur A. pinnala bark thin, dull gray brown, smooth then fissured with age. Leaves brilliant orange i Maple A. palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from Maple A. palmatum small slow growing, in autumn gold orange and red. Underplant to shelter from Maple A. palmatum small slow growing, small tree. Yellow leaves in spring, in autumn the dissectum Deeply cut lace-like purple leaves. 2m high with broad, drooping habit Maple A. shirasawanum Aureum slow growing small tree. Yellow leaves in spring, in autumn ma Snakebark A. capillipes 10–15m often multiple trunks. Bark olive green, grey stripes. Male/female Sugar A. saccharinum 45m. Leaves bright yellow to orange to fluorescent red-orange. Flowers earb Sycamor A. pseudoplatamum bellitantissimum' slow growing intense salmon pink leaves, later yello lar Common Nottingham Mespilus germanica Russett apple like fruit requires bletting to cat (see beary Black Morus nigar 10–15m. Self-fertile, dark purple fruit. SW Asia, 154 pairs of chromosom Quercus beech family, Fagaseas. Spirally arranged leaves, acoms contain tannic acons,	-like fruits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour.  female on separate plants, wind polilianced. Berry-like seeds, taking 18 months to ripen.  nd fall in late autumn. cones are creet, small, green or purple, ripening brown 5-8 months after pollination, northe grain for modelling, carving  t small, round, ribbed eream audet  somewhat toxic to bumblebees.  a much visited by bees. Fruit oval  eaves, mostly tolerating limestone soils and wind.  to red in autumn.  nd driving winds and sun. To 5m.  Winter bank attenctive. 2m arching habit  e cream areas turn deep searlet  ungined red, strong colour from good sun  mental effect. / to control size  unumn. Red flowers in spring.  flowers on different racemes.  y spring.  w green. Monoecious.  satiring to deexy and ferment). Ornamental and hardy, slow and graceful in growth. White flowers in June.  ses. Branches pruned affer leaves fall can be used to make baskets. White sap in raw fruit & green parts intoxicating all female flowers in spring.  flor rucky places?.  nd purple in auturn  at after for furniture making.  litin A  Sanguinole 2, Aug/Sep. Small, russec, red blush & spotted. Flesh red stained, juicy and musky QA  ribens  ribens Bon Chretten 2,12 Aug/Sep.Fat fresh Ripe yellow QA  Wither Nellis  saller, Perus Communis, Wick Coart  Judge Amphlett Rec, 1, late September, 1 week, med A, low T, avg Q  Moorcroft 2, late Sept, N 2d, med A, med T, avg to good Q  Red Pear 5, Oct. >3 wks, lowA, lowT, avg Q  Snake Pole From Adnigham  Soneyway N/a  Sweet Huffcap=Helens Early 3, Sept, 1 wk, med A, low-med T, avg Q  Thorny Alars lept, 1 week, med A, low T, avg Loy good Q  Thurston's Red 3, 2/3wk Oct, M<3wks, med A, low-med T, very pood Q  Velow Huffcap 2, early with Oct, 1 week, med-hi A, low'T, fruit v.g. Q  Snake Pole From Adnigham  Sone plant Prom Adnigham  Sone plant Prom Adnigham  Sone pine P. pinea 20m cones 36 months to mature. Ptdbble nuts  sto vinvikel-Vernis mid summer.  k mature in summer  k mature in summ
per Juriperis communis foliage of the juvenile needle-like type, unlike pine. Dioccious, male and I. Larix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, nurn yellow are as have small cones with short bracts.  2 Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little & Proat-leaved Lime Tpataphyllos vellow-white flowers (make lime tea) pollinated by boes, frui a Silver lime T. Communiosa flowers and event pollinated by horse-free should be supported by the pollinated by boes, frui a Silver lime T. Contact flowers are received by hermaphrodite, pollinated by horse-free should be supported by the pine supported by the pollinate of the supported by the pine suppor	Hille Fraits dark-red when young, but become dark-purple when ripe, edible and have an apple and sweet flavour. Fernale on separate plants, wind polinated. Berry-like seeds, tailing 18 months to repen. and fall in late autumn. cones are erect, small, green or purple, ripening brown 5-8 months after pollination; northe grain for modelling, carving.  1 small, round, ribbed cream nutlet.  1 somewhat toxic to bumblebees.  2 much visited by bees. Fruit owal concesses are controlled to the state of t
per Juriperis communis foliage of the juvenile needle-like type, unlike pine. Dioccious, male and I. Larix decidua 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are as have small cones with short bracts.  2 Tilia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little A Broad-leaved Lime Tpataphyllos yellow-white flowers (make lime tea) pollinated by bees, frui 3 silver lime Ttomcruson flowers pale vellow, hermaphrodite, pollinated by honeybees. Nectar is 5 small-leaved lime Teordatat flowers grone-yellow, hermaphrodite, pollinated by honeybees. Nectar is 5 small-leaved lime Teordatat flowers grone-yellow, hermaphrodite, pollinated by honeybees. Nectar is 5 small-leaved lime Teordatat flowers grone-yellow, hermaphrodite, pollinated by honeybees. Nectar is 5 small-leaved lime Teordatat flowers grone-yellow, hermaphrodite, pollinated by honeybees. Nectar is 5 small-leaved lime Teordatat flowers groneybee and the flowers of the flowers flowers and the proper state. The Translation of the flowers of the flowers of the flowers flowers. The translation of the flowers flower	-like Fraits deals red when voung, but become dark-purple when rine, edible and have an apple and sween flavour. Fernation on sparnet plants, wind pollinated. Berry, like seeds, taking 18 mouths to ripen.  Ind fall in lace autumn. cones are creet, small, green or purple, ripening brown 5.8 months after pollination; norther grain for modelling, carving it small, round, ribbed cream nutlet.  Somewhat toxic to burblichees.  In much visited by bees. Frait toxal carves, mostly role arising limestone soils and wind.  To red in autumn.  In drung winds and sun. To 5m.  Winter burk structive. 2m arching habit  ceream areas turn deep scalet  regined red, strong colour from good sun montal effect. / to control size turnum. Red Bowes in spring.  flowers on different racenes.  Spring.  In write modes and ferment; Ornamental and hardy, slow and graceful in growth. White flowers in June.  Ess. Branches pranted after leaves fall can be used to make baskets. White sap in raw fruit & green parts intoxicating half from a for furniture making.  Jilin A  Sanguinole 2, Aug/Sep. Small, russet, red blush & spotted. Flesh red stained, juicy and musty QA  Thompson's Sep, store to Oet or Nov. Golden skin, patch of russet, Flesh hisiry with rich performe Williams Bon Chretien 2,12 Aug/Sep.Etat fresh Ripe vellow QA  Winter Nellis  saller, Pyrns. Communis, Wick Court  Judge Amphlert Rec, I, late September, I week, med A, low T, avg Q  Moorcroft 2, late Sept, Mad, med A, med T, savingen good Q  Red Pears, S. Cep. 35 wks, JuwA, John, J. avg to good Q  Red Pears, S. Cep. 35 wks, JuwA, John, J. avg to good Q  Thurston's Red 3, 2/3wk Oet, McSwks, med A, Jow T, avg Q  Volloried G., Oet, 3-6 wks, med A, Jow T, avg to good Q  Vollow Huffeap P-Lelens Early 3, Sept, I wk, med A, Jow T, good Q  Villow Huffeap P-Lelens Early 3, Sept, I wk, med A, Jow T, good Q  Vollow Huffeap P-Lelens Early 3, Sept, I wk, med A, Jow T, fruity v.g. Q  Shake Pole From Oventon, very old general purpose  Speatt From Affalbam  Stoneyway N/a  Sweet Huffeap P-Lelens Early 3,
per Juniperis communis foliage of the juvenile needle-like type, unlike pine. Dioecious, male and in Luxir decidual 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow as cas have small cones with short braces.  **Titla sturdy trunk, branches divide into numerous ramifications.**Timber soft, easily worked, little A Broad-leaved Lime **Lingathyribios* yellow-white flowers (make lime tea) pollinated by bose, freil a Silver lime **Lingathyribios* yellow-white flowers (make lime tea) pollinated by boneybees. Nectar is a Silver lime **Lingathyribios* yellow, hermaphrodite, pollinated by honeybees. Nectar is Small-leaved lime **Lioyabacaa.**Sturdy widely branched deciduous tree with bandsome palmate for hum **Lingathyribios* yellow, hermaphrodite, pollinated by honeybees. Nectar is small the human spirate in a stade when yellow have spirated with a gate. Leaves brilliant orange. Maple **A. palmatum* small slow growing, in autumn gold orange and red. **Chelephant to shelter from **Lindahe** after grower's genal daughter. Decply cut leaves vividly marked with pink and white. **Maple **A. palmatum* small slow growing, in autumn gold orange and red. **Chelephant to shelter from **Lindahe** after grower's genal daughter. Decply cut leaves vividly marked with pink and white days the state of the palmatum* shelt grower is grower and the state of the	Like fruits dark sed when young, but become dark purple when ripe, edible and have an apple and sweet flavour. Formals on separate planes, wind polinated. Berry-files excels, taking its months to ripon.  Ind full in late autumn, cones are erect, small, green or purple, ripening brown 5-8 months after pollination; northe grain for modelling, carving it small, round, ribbed eream nutlet is somewhat took to bumblebees, growth wished by bees, Frain oral excess, mostly obsided took took bumblebees, growth wished by bees, Frain oral excess, mostly obsided and sum. To 5m.  Winter bark attractive. 2m arching habit excess mostly obsided and sum. To 5m.  Winter bark attractive. 2m arching habit excess and excess and experiment deep scadet experiment of the common and the product of the common and common and the com
per Juniperis communis foliage of the juvenile needle-like typer, unlike pine. Dioecious, male and in Laris decidual 15-50 m leaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small cones with short bracts.  **Piña strucky trunk, branches divide into numerous ramifications.**Timber soft, easily worked, little **M Broad-leaved Lime**Diaphyllos yellow-white flowers (make lime tea) pollinated by bees, fruit a Siliver lime **T.tomentosa** flowers pace vellow, hermaphrodite, pollinated by honerbees. Nectas it Small-leaved lime **T.comentosa** flowers pace vellow, hermaphrodite, pollinated by honerbees. Nectas it Small-leaved lime **T.comentosa** flowers pace vellow, hermaphrodite, pollinated by honerbees. Nectas it Small-leaved lime **T.comentosa** flowers powers pand-daughter. Deeply cut leaves vividly marked with pink and white. Mapte **A.p.fimatah** flat for grower's grand-daughter. Deeply cut leaves vividly marked with pink and white. **Mapte **A.p.fimatah** flate grower's grand-daughter. Deeply cut leaves vividly marked with pink and white **Mapte **A.p.fimatah** flate grower's grand-daughter. Deeply cut leaves vividly marked with pink and white **Mapte **A.p.fimatah** flate grower's grand-daughter. Deeply cut leaves vividly marked with pink and white **Mapte **A.p.fimatah** flate grower's grand-daughter. Deeply cut leaves vividly marked with pink and with pink in autumn the disease time been been grower flate growers and grower flate growers	Like Fruits dark red when young, but become dark purple when ringe, edible and have an apple and sweet flavour. Email or negratine plants, wind pollimated. Berr-like seeds, baking H months to ripem.  Ind fill in lare autumn, cones are erect, small, green or purple, ripening brown 5-8 months after pollination, norther grain for modelling, carving.  It mall, round, Shibed cream mulet  Counceshart tracic to bumblebees,  Journal white discovering investments with a contract of the state of the state of the contract of the state of the s
per Juniperis communis foliage of the invende needle like type, unlike pine. Dioccious, male and a Larix decidents 15 90 m leves needle like, borne singly spirally on the long shoots, rum yellow are shave small comes with short bracts.  Filia sturdy trunk, branches divide into numerous ramifications. Timber soft, easily worked, little & Broad-leaved Lime Tejharyph/filos yellow-white flowers (make lime teal pollinated by honerbees. Nectas is 3 silver lime. Tetomenosa flowers pale vyllow, hermaphrodite, pollinated by honerbees. Nectas is 5 small-leaved lime. Tecordated flowers pene-yellow, hermaphrodite, ofbliod in salads when you gowing to Acer Of the family Apindaceae. Sturdy widely branched deciduous tree with bandsome palmate is 5 small-leaved lime. Tecordated flowers pene-yellow, hermaphrodite, ofbliod in salads when your large. Apalmatum small slow geowing, in autumn bed orange and red. Underplant to shelter fore. [Ariation*] firet growers, grand-daughter. Deeply cut leaves vividly marked with pink and white flowers in June Prune hard in spring, in autumn the dissectum Deeply cut lace-like purple leaves. 2m light with broad, decoping habit advanced with pink and with the plant of the dissectum Deeply cut lace-like purple leaves. 2m light with broad, decoping habit and plant of the	-like fruits dark-red when young, but become dark-pupels when ripe, edible and have an apple and sweet flavour. female on separate planes, wind polithard. Berny-flies ecols, taking its months origin.  dial in lac autumn, cones, are exect, small, green or pupple, riponing brown 5-8 months after pollination; norther grain for modelling, carving it small, round, blobed eream nurbet is sumewhat tooks to bumblebees. A meet and the sumewhat tooks to bumblebees. The problem of the sumewhat tooks to bumblebees and the sumewhat tooks to bumblebees. The problem of the sumewhat tooks to bumblebees. The problem of the sumewhat tooks to bumblebees and the sumewhat tooks to bumblebees. The problem of the sumewhat tooks to bumblebees and the sumewhat tooks and sum. To fine whether the sumewhat tooks are summed after 1 to fine. The sumewhat tooks are summed after 1 to someth size until the sumewhat tooks and the sumewhat tooks are summed after 1 to someth size until the sumewhat after 1 to someth size until the sumewhat the summer of the
per Juniperis communis foliage of the inventie needle like type, unlike piace Dioccious, male and a Laris decidual 15-30 in leaves needle-like, borne singly spirally on the long shoots, turn yellow are shave small comes with shurt brachs.  6 Tilia sturds trunk, branches divide into numcrous ramifications. Timber soft, easily worked, little A Broad-leaved Lime T. plaryphytlos yellow white flowers (make lime tea) pollinated by becas, fetti 8 Silver lime T. turnemiosa Howers peak vellow, hermaphradite, pollinated by boneshees. Necroit 9 Silver lime T. turnemiosa Howers peak vellow, hermaphradite, pollinated by boneshees. Necroit 9 Silver lime T. turnemiosa Howers peak vellow, hermaphradite, pollinated by boneshees. Necroit 9 Silver lime T. turnemiosa Howers peak vellow, hermaphradite, pollinated by honeshees. Necroit 9 Similal 16 Acer Of the family Symbolosose. Sturdy widely branched deciduous tree with bandsome palmate to Ammur A. Jamala bark thin, dull graw brown, smooth them fissured with a gue a substance of the similar shadows proving, in autumn gold orange and red. Underplant to shelter from the dissectum Deeply cut lace-like peak of the complex of the similar shadows and the	-like frais dark-red when vooras, but become dark-purple when rice, edible and have an apple and sweer flavour. female on separate plans, with pollimation; norther formation and fall in late autumn, comes are erect, small, green or purple, apening brown 5-8 months after pollimation; norther gram for modelling, carving it is mall, round, photo decam outlet a somewhat took to bumbibless.  yourch where the beste decam outlet a somewhat took to bumbibless.  yourch where the beste decam outlet a somewhat took to bumbibless.  yourch where the beste 1-not road average and to the somewhat took to bumbibless.  where the strained outlet is the somewhat took to bumbibless.  or cam areas num.  In drained when your decame to the somewhat to bumbible to cram areas num deep scarlet agened text, strong colour from good summental effect of use outlets are summented flowers in spring.  Bowest out different recemes: typing.  Bowest out different recemes.  Typing.  we green. Monocolous.  textens to decar and ferment). Ornamental and hardy, slow and graceful in growth. White thowers in June.  The Branches pruned after leaves fall can be used to make baskets. White sup an new fruit & green parts intoxicuting all female flowers in spring.  Yound, lower once droop  of order places.  A Sunguinole 2, Aug/Sep. Small, russet, red blosh & spotted. Flesh red stained, jucy and mosky QA deleas.  A Sunguinole 2, Aug/Sep. Small, russet, red blosh & spotted. Flesh red stained, jucy and mosky QA deleas.  A Sunguinole 2, Aug/Sep. Small, russet, red blosh & spotted. Flesh red stained, jucy and mosky QA deleas.  A Sunguinole 2, Aug/Sep. Small, russet, red blosh & spotted. Flesh red stained, jucy and mosky QA deleas.  A Sunguinole 2, Aug/Sep. Small, russet, red blosh & spotted. Flesh red stained, jucy and mosky QA deleas.  A Sunguinole 2, Aug/Sep. Small, russet, red blosh & spotted. Flesh red stained, jucy and mosky QA deleas.  A Sunguinole 2, Aug/Sep. Small, russet, red blosh & spotted. Flesh red stained, jucy and mosky QA deleas.  A Sunguinole 2, A

Black Walnut J.nigra large fast-growing with distinctive ridged bark. Likes most soil types, deep sites, not liking late frosts. To 25m Common Walnut *J.regia* 30m slow-growing, round-headed native, foliage very aromatic when young. Nuts taste better picked unripe.

Cricket Bat Willow S. alba 'Caerulea' 10-20m. Dioecious. Used for cricket bats, a tough, lightweight wood that does not splinter easily. Purple Willow S.purpurea To 5m. Purple catkins. purple-brown-yellow shoots, pale grey on old stems. Also used to make cricket bats

Sallow or Goat Willow S.caprea grows in any soil, wet or dry. Prized for its silky catkins with yellow anthers in March, valuable for bees. Scarlet Willow S.alba Chermesina/Britzensis wet or dry soil. Waxy yellow, orange and red young stems, usually coppiced for these. Weeping Willow S.sepulcralis 'Chrysocoma' To 20m. Shoots and foliage gold in spring, green in summer, autumn gold again.

Yezo Dwarf Willow S.nakamurana var yezoalpina low growing, prostrate species of willow with lovely catkins which can be found growing in the wild on the alpine slopes of Hokkaido, Japan

Corkscrew Willow S.babylonica tortuosa young shoots used in flower arrangements, putting out roots. Dioecious. To 10m Crack Willow S.fragilis bark grey-brown leaves bright green, serrated margin. Catkins early spring pollinated by insects.